Optimal Fresh

The fruit, vegetable and fresh produce expert system



Detailed Report Printed on Tuesday, 18 December 2001

Crop apricot

Maturity stage General
Category Fruit
Plant Part Fruit

Usage Baked, Fresh/ Raw,

Juice/ Drink, Processed/ Canned

Botanical name Prunus armeniaca L.

Botanical family Rosaceae



Picture source: Sydney Postharvest Laboratory, 1999

Alternate names include

Refrigerated Container/Coolroom Recommendations

Optimum product storage temperature

-0.5 to 0.0°C

0.0°C

Temperature set point

Add a margin for uncertainty in equipment performance if necessary. For return air control set point add 1°C to delivery set point.

Ventilation (air exchange) settings for containers: $6 \text{ m} (20^\circ) = 60 \text{ m}^3/\text{h} = 35 \text{ cfm}$

12 m (40') = **120 m³/h = 70 cfm**

Acceptable product temperature at loading into container

-0.5 to 4.5°C

Key Properties

Storage time	Humidity	Freezing point	Storage time at	Ventilation
(days)†	(% RH)	(°C)	ambient (~20°C)	rate
14 - 21	90 - 95	-1.1	1 - 2	High

[†] at optimum storage temperature

Depends on cultivar

Other Properties

Ref	Maturity stage	Air exchange *	Freezing Point (°C)	Ethylene production **	Ethylene sensitivity	Ice compat- ibility	Water loss ***	% Water content	Bruising suscept- ibility
1	General	High	-1.1	High	High	No	Η	85.4	

^{*} Air exchange rates: Nil = 0%; Very low = 25%; Low = 50%; Medium = 100%; High = 200%; Very high = 400% fresh air/hour.

^{**} Ethylene production rates at 20°C: Nil = 0 nM; Very low = <4 nM; Low = 4 - 40 nM; Medium = 40 - 400 nM; High = 400 - 4000 nM; Very high =>4000 nM ethylene/kg/hour.

^{***} Where % weight loss/week is given this is converted as: Low <= 1%; Medium = 1.1 - 3.4%; High = >3.5%

Optimal Fresh

The fruit, vegetable and fresh produce expert system



Controlled Atmosphere

Ref	Maturity stage	% O2		% CO2		Temp°C		Benefit of controlled	
		min	max	min	max	min	max	atmosphere	
1	General	2	3	2	3	0	5	Slight, (+0 days)	

Reference notes

1 CA not recommended for fresh-market apricots

Respiration* and Heat Transfer

Ref Maturity		0	°C	5	°C	10)°C	15	O°S	20)°C	25	s°C	Specific heat
	stage	min	max	kJ/kg/EC **										
1	General	16	17	33	56	63	102	88	156	136	200			3.70

^{*} Respiration values given are in Watts per tonne. 1 W/t = 20.4 kCal/t/d = 82.1 Btu/tn./d = 73.3 Btu/2000 lbs/d = 0.167 mL CO2/kg/h = 7.0 umol CO2/kg/h = 0.308 mg CO2/kg/h

Compatibility in Mixed Storage

Temperature compatibility group

0	7	13	20
	-		

Humidity compatibility group

Dry	Moderate	High	Very high
60-80%	80-90%	90-95%	95-100%

Not compatible with crops that: Produce ethylene (especially when they are ripe or ripening)

Are sensitive to ethylene

Odours will be absorbed by:

Absorbs odours from:

Ethylene-producing fruits and vegetables from Optimal Fresh database

(Medium ethylene production levels or greater.)

apple	apricot	atemoya	avocado
banana	breadfruit	cherimoya	custard apple
durian	feijoa	fig	jackfruit
jujube fruit	kiwifruit	litchi	mamey sapote
mango	mangosteen	melon, cantaloupe	melon, honeydew passionfruit
nashi	nectarine	papaya	
peach sapodilla	pear tomato	plum	rambutan

^{**} Specific heat (kJ/kg/°C) = 0.0335 x % water content + 0.8374; Specific heat in Btu/lb/°F = 0.08 x % water content + 0.2

Optimal Fresh

The fruit, vegetable and fresh produce expert system



Ethylene-sensitive fruits and vegetables from Optimal Fresh database

(High sensitivity.) Chinese cabbage apricot Chinese broccoli apple banana asparagus atemoya avocado bean, French bitter melon bok choy broccoli cabbage cauliflower brussels sprouts carrot cherimoya celery chicory collards corn, sweet cucumber custard apple eggplant endive fuzzy melon globe artichoke guava kale kiwifruit kohlrabi leafy greens lettuce litchi lemon long bean mandarin mangosteen mamey sapote mango melon, cantaloupe melon, honeydew nashi nectarine olive, fresh okra onion, green papaya parsnip passionfruit pea, green peach pear persimmon plum potato pumpkin quince rambutan rhubarb sapodilla silver beet spinach squash, soft rind squash, zucchini sweet potato tamarillo tomato turnip greens watermelon yam

Seasonal Availability

Ref	Country	Region (where given)	Start Season	End Season	Start Peak	End Peak
1	Canada		July	August	-	-
1	Chile		November	December	-	-
1	Australia		October	March	November	January
1	UK		July	August	-	-
1	USA		January	August	-	-

References for apricot

Values quoted in Detailed Report are taken from a compilation of the best set of figures from all references. This best set of figures is always referred to as Reference 1.

See Reference Report for full listing of all values, original references and alternate names.